

Formiminotransferase Cyclodeaminase (LC1)

Antigen Specification

Product Number: 13700

Description:

Human formiminotransferase cyclodeaminase (FTCD); identical with the antigen called "liver cytosol type 1 antigen" (LC1). Recombinant antigen for in vitro research and manufacturing use only.

Immunological function:

Binds IgG-type human auto-antibodies.

Origin:

Recombinant. Expressed by recombinant baculovirus (*Autographa californica* multiple nuclear polyhedrosis virus; AcMNPV) infection of *Spodoptera frugiperda* Sf9 insect cells.

Expression construct:

Full-length cDNA coding for the approx. 60 kDa monomer chain of the human FTCD enzyme fused to a hexahistidine purification tag.

Biochemical tests:

SDS-PAGE (purity > 95 %); Western blot with i: monoclonal anti-His-tag antibody; ii: anti-LC1 autoantibodypositive sample.

Calculated isoelectric point:

pH 5.85

Calculated molecular weight: 60 kDa

(The native human enzyme has been reported as an octamer with approx. 480 kDa molecular weight.)

Immunological tests/Functionality:

Standard ELISA test (checkerboard analysis of positive/negative samples); immunodot analyses with positive/negative samples.

Recommended buffer/storage and handling conditions:

Recommendations for storage buffer: neutral to slightly alkaline pH and 20 % glycerol as cryoprotective agent. Storage conditions: –70° or below. Repeated freeze/thaw cycles should be avoided.

Coating concentration:

0.35-0.7 µg/mL (depending on the type of ELISA plate and coating buffer; <u>CAUTION</u>: these values are not based on exhaustive testing and are only guidelines for customer's development work). Suitable for labeling of functional groups.

Copyright 2001 - 2021 DIARECT GmbH

DIARECT GmbH • Bötzinger Str. 29 B • 79111 Freiburg • Germany • Tel. +49 (0) 761 47979-0 • Fax +49 (0) 761 47979-29 info-dia@bbisolutions.com • www.bbisolutions.com Antigen Specification LC1_13700_210421_Rev05.doc